arXiv:1303.7406v1 [math.DG] 29 Mar 2013

PRESCRIBING METRICS ON THE BOUNDARY OF CONVEX CORES OF GLOBALLY HYPERBOLIC MAXIMAL COMPACT ADS 3-MANIFOLDS

BOUBACAR DIALLO

Abstract. We consider globally hyperbolic maximal anti de Sitter 3-manifolds M with a closed Cauchy surface S of genus greater than one and prove that any pair of hyperbolic metrics on S can be realized as the boundary metrics of the convex core of a maximal globally hyperbolic anti de Sitter 3-manifold structure on M. This answers the existence part of a question of Mess about the unique realization of such metrics. Our theorem has a nice formulation purely in terms of 2-dimensional Teichmüller theory and earthquakes.

Contents

1. Introduction 2
1.1. Globally hyperbolic maximal compact Anti de Sitter manifolds 2
1.2. Analogies with quasifuchsian hyperbolic 3-manifolds 3
1.3. The Mess conjecture 3
1.4. Relation to Teichmüller theory and earthquakes 3
2. Basic results and strategy of the proof 5
2.1. Anti de Sitter geometry in dimension 3 and globally hyperbolic manifolds 5
2.2. How to show that Φ is onto 5
Acknowledgements 6
3. Outline of the proof of the properness theorem 6
4. Proof of the main propositions 8
5. Computing degrees 14
References 16

Date: April 1, 2013.

partially supported by the A.N.R. through project ETTT, ANR-09-BLAN-0116-01, 2009-13.

1. Introduction

1.1. Globally hyperbolic maximal compact Anti de Sitter manifolds.

Anti de Sitter space (AdS) is the model space of Lorentzian manifolds of constant sectional curvature equal to -1. Its $n-$dimensional avatar can be defined as the -1 level set of the standard nondegenerate quadratic form of index 2 in $\mathbb{R}^{n+1}$, with the induced quadratic form on each affine tangent subspace (it has index one on such subspaces).

An anti de Sitter 3-manifold is thus a (smooth, connected) manifold M endowed with a symmetric bilinear covariant 2-tensor of index one, everywhere nondegenerate, whose sectional curvatures are all equal to -1. By classical results, such an M is locally isometric to the model space AdS_3, anti de Sitter space of dimension 3. If we restrict ourselves to oriented and time oriented manifolds, it is therefore endowed with a (G, X) structure, where X is AdS_3 and G the identity component of its isometry group. Both definition are equivalent in that case. Let's call such objects AdS_3 spacetimes.

An AdS_3 spacetime is said to be globally hyperbolic if it admits a Cauchy hypersurface: a spacelike surface which intersect every inextendable timelike line exactly once. If the spacetime has a compact Cauchy surface, then every Cauchy surface is compact. Moreover if the spacetime cannot be isometrically embedded in a stricty larger spacetime by an isometry sending a Cauchy surface to another one, then it is called globally hyperbolic maximal compact (GHMC).

(G, X) manifolds (with G and X as above) have a well defined holonomy representation, up to conjugation, and a developing map (up to equivalence).

Recall that the identity component of the isometry group of AdS_3 identifies (up to index 2) with $PSL_2(\mathbb{R}) \times PSL_2(\mathbb{R})$. Thus holonomies of (G, X) spacetimes M are represented by pairs of representations from the fundamental group of M to $PSL_2(\mathbb{R})$. In the case where M is globally hyperbolic maximal with a surface S of negative Euler characteristic as a Cauchy surface, one of Mess's main theorem ([12, proposition 19], see also [1]) asserts that such a pair (ρ_l, ρ_r) is a point of $\mathcal{T}(S) \times \mathcal{T}(S)$, the product of two copies of the Teichmüller space of S. Recall that $\mathcal{T}(S)$ is the space of discrete and faithful representations of $\Pi_1(S, x)$ (with any choice of basepoint x) to $PSL_2(\mathbb{R})$ modulo conjugation. Equivalently, it is the space of metrics on S of constant sectional curvature -1, modulo the equivalence relation identifying two of them iff there is a diffeomorphism isotopic to the identity, which is an isometry from one metric to the other.

1.2. **Analogies with quasifuchsian hyperbolic 3-manifolds.** Mess noted that his theorem on holonomies of $GHMC$ anti de Sitter spacetimes of dimension 3 is the analog of the simultaneous uniformization theorem of Bers for quasifuchsian hyperbolic three manifolds [3]. Thurston asked whether one could uniquely prescribe the two hyperbolic metrics on the boundary of the convex core of such a manifold [15], [6]. So far, only existence has been proved, thanks to works of Epstein and Marden [7] on Thurston and Sullivan's $K = 2$ conjecture (which happens to be false, see [8]), and Labourie [10], independently.

In his work [12], Mess established further analogies between quasifuchsian hyperbolic 3-manifolds and $GHMC$ AdS_3 spacetimes. Indeed such spacetimes M have a well defined convex core, which as in the hyperbolic setting is the minimal non-empty closed convex subset. Except for the Fuchsian case where both upper and lower boundary metrics are equal, the convex core has two boundary components which are pleated hyperbolic surfaces. Both are thus equipped with hyperbolic metrics and measured bending laminations. In particular this defines a map Φ from the space of $GHMC$ AdS structures (with Cauchy surface S of fixed topological type), identified with $\mathcal{T}(S) \times \mathcal{T}(S)$, to $\mathcal{T}(S) \times \mathcal{T}(S)$, which sends a structure to the ordered pair of upper and lower boundary hyperbolic metrics. In the quasifuchsian setting, the analogous map is then onto.

1.3. **The Mess conjecture.** Mess asked whether the map Φ is one-to-one and onto, that is, whether an ordered pair of hyperbolic metrics on S can be uniquely realized as the upper and lower boundary metrics of a $GHMC$ AdS_3 spacetime M. This is the analog of Thurston's conjecture for quasifuchsian manifolds. Uniqueness is still an open question. The present article gives a positive answer to the existence part of this conjecture of Mess.

Note that our statement, in the anti de Sitter setting, cannot be proved by methods of Epstein and Marden. Indeed, the K-quasiconformal constant in their theorem cannot exist in our context, because it would contradict the earthquake theorem (see next section). There's no restriction on the bending measures of our spacetimes, as opposed to the hyperbolic setting. Moreover, the analog of Labourie's theorem [10, theorème 1] or of Labourie and Schlenker's theorem [11] remains unknown in the AdS setting.

1.4. **Relation to Teichmüller theory and earthquakes.** Earthquakes were defined by Thurston in [16] by extension to measured laminations of the case of simple closed curved. Let E^l_λ and E^r_λ be the left and right earthquakes along a measured lamination λ. Thurston proved that those two maps from $\mathcal{T}(S)$ to itself are (continuous and) bijective and in fact are inverse to each other. His earthquake theorem asserts that for any m and m' in $\mathcal{T}(S)$, there is a unique measured lamination λ (resp. λ') such that the left (resp. right) earthquake along λ (resp. λ') sends m to m'.

Mess rephrased this earthquake theorem in the context of pleated surfaces in AdS_3 geometry.

Let S be a closed surface of negative Euler characteristic, M a globally hyperbolic maximal AdS_3 spacetime with S as a Cauchy surface, (ρ_l,ρ_r) the holonomy of M. Let λ_+, λ_- the upper and lower pleating laminations of the convex core of M, m_+ and m_- the corresponding boundary hyperbolic structures.

Then

$$\begin{aligned}\rho_l &= E^l_{\lambda_+}(m_+)\\ \rho_r &= E^r_{\lambda_+}(m_+)\\ \rho_l &= E^r_{\lambda_-}(m_-)\\ \rho_r &= E^l_{\lambda_-}(m_-)\ .\end{aligned}$$

$$\begin{tikzcd}
 & m_+ \arrow[dr, "E^l_{\lambda_+}"] & \\
\rho_l \arrow[ur, "E^l_{\lambda_+}"] & & \rho_r \arrow[dl, "E^l_{\lambda_-}"] \\
 & m_- \arrow[ul, "E^l_{\lambda_-}"] &
\end{tikzcd}$$

Figure 1. Mess diagram

Thanks to Thurston's theorem, our map Φ of the previous section is continuous. Via the earthquake theorem, surjectivity of Φ is thus equivalent to the following statement:

Prescribing middle points of 2 intersecting earthquakes paths For any two points m_+ and m_- in $\mathcal{T}(S)$, there are left and right earthquakes, joining two points say ρ_l and ρ_r in $\mathcal{T}(S)$, such that m_+ and m_- are the middle points of the corresponding earthquake paths.

Again we fail to prove uniqueness.

2. Basic results and strategy of the proof

2.1. **Anti de Sitter geometry in dimension 3 and globally hyperbolic manifolds.** Anti de Sitter 3-space is the model space of Lorentzian 3-manifolds of constant sectional curvature equal to -1. Like hyperbolic space, it has a boundary at infinity, Einstein space of dimension 2, with a conformal Lorentzian structure, and it has several equivalent definitions (which give isometric spaces).

Apart from the quadric model we already mentioned, in dimension 3, it is also the Lie group $SL_2\mathbb{R}$ with its induced Killing metric. The Lie group model allows us to identify the group of isometries of AdS_3 with

$$SL_2\mathbb{R} \times SL_2\mathbb{R}/J \ ,$$

where J is the subgroup of order 2 generated by the matrix couple $(I_2, -I_2)$. Therefore this isometry group is a double cover of $PSL_2\mathbb{R} \times PSL_2\mathbb{R}$. The space of holonomies of AdS_3 spacetimes (modulo conjugation) is thus a double cover of the space of representations of $\pi_1(M)$ to $PSL_2\mathbb{R} \times PSL_2\mathbb{R}$ (modulo conjugation). A point in that representation space will be called a pair of left and right holonomy.

A theorem of Mess asserts that, for a GHMC AdS_3 spacetime M, with Cauchy surface S, the left and right holonomies are both discrete, faithful representations which act (properly discontinuously and) cocompacty on the hyperbolic plane: these are Fuchsian representations. Conversely Mess proved that to such a pair of Fuchsian representations is the holonomy pair of a unique GHMC AdS_3 manifold M homeomorphic to $S \times \mathbb{R}$ (in particular, $\pi_1(x, M)$ and $\pi_1(x, S)$, with suitable choice of basepoint x, are isomorphic). This is the exact analog of Bers theorem for quasifuchsian hyperbolic 3-manifold. This gives a natural homeomorphism between the space of GHMC AdS_3 structures (with orientation and time orientation) and $\mathcal{T}(S) \times \mathcal{T}(S)$.

2.2. **How to show that Φ is onto.** Recall that Φ is the map from $\mathcal{T}(S) \times \mathcal{T}(S)$ to itself which associates to a GHMC structure on a manifold M with $\pi_1(M) \cong \pi_1(S)$ the ordered pair of upper and lower boundary metrics on its convex core. Thanks to the earthquake theorem, Φ is continuous. We just need to show that it is a proper map, and since it will have a well defined degree, that it is a map of degree one. It then follows that ϕ is onto.

Properness theorem:

With notations as in the previous sections, if m_+ lies in a compact set in $\mathcal{T}(S)$, m_+ and m_- lie in the image of Φ, if preimages ρ_l and ρ_r in $\mathcal{T}(S)$ are such that ρ_l leaves every compact subset of $\mathcal{T}(S)$, then so does m_-.

We may rephrase it with sequences in $\mathcal{T}(S)$ instead of compact sets:

if $(m_+^n)_{n\geq 0}$ is a sequence in $\mathcal{T}(S)$ converging to m_+^∞ in $\mathcal{T}(S)$, $(m_-^n)_{n\geq 0}$, $(\rho_l^n)_{n\geq 0}$ and $(\rho_r^n)_{n\geq 0}$ are sequences in $\mathcal{T}(S)$ such that $(\rho_r^n)_{n\geq 0}$ tends to ∞, then $(m_-^n)_{n\geq 0}$ also tends to ∞.

In our context, we can (equivalently) replace "'$(\rho_r^n)_{n\geq 0}$ tends to $(\infty$" by "'$l_{m_+^n}(\lambda_+^n)$) tends to ∞". See next section for further details.

Degree theorem:

Φ is properly homotopic to a homeomorphism. Since a homeomorphism has degree one, it then follows that our map Φ has degree one.

Acknowledgements

I'd like to thank Jean Marc Schlenker for his grateful help and guidance during this work. I would also like to thank F.Bonsante, T.Barbot, C.Series, S.Kerckhoff, J.Danciger for fruitful discussions.

3. Outline of the proof of the properness theorem

Our main result here is the following theorem:

Theorem 1 (properness of Φ). *Let Φ be the map defined in the previous section. Then Φ is a proper map.*

We prove this theorem with 3 propositions, which will be proven in the next section.

The first one is a slight refinement of Lemma 4.2 of [5].

Let S be a closed surface of negative Euler characteristic, $\alpha_0 > 0$. Let $g \in \mathcal{T}(S)$ and let c a closed geodesic for g. For any point $x \in c$, let $g_x^r(\alpha_0)$ (resp. $g_x^l(\alpha_0)$) be the geodesic segment of length α_0 (for the metric g), orthogonal at x to the right (resp. the left) of c. Let $n_l(x)$ (resp. $n_r(x)$) be the intersection number of $g_x^r(\alpha_0)$ (resp. $g_x^l(\alpha_0)$) with c, including x.

Proposition 2. *There exists a constant l_0 (depending only on α_0 and the genus of S) such that for all $\delta_0 > 0$, there exists some $\beta_0 > 0$ (depending on α_0, δ_0 and the genus of S) such that:*

$$l_g(\{x \in c : \inf(n_l(x), n_r(x)) \leq \beta_0 l_g(c)\}) \leq \delta_0 l_g(c) + l_0.$$

What the proposition says is: if the length of c is large enough, for most points x of c, the left (resp. right) going arc orthogonal to c at x, of fixed length α_0 intersects c a lot.

By density of weighted multicurves in $\mathcal{ML}(S)$ and continuity of length functions [9] and intersection numbers, we get a similar version for more general measured laminations on S, namely:

Corollary 3. *With the same notations as above, let $\lambda \in \mathcal{ML}(S)$ be a measured lamination whose support is a simple closed curve c. Then there exists a constant l_0 (depending only on α_0 and the genus of S) such that for all $\delta_0 > 0$, there exists some $\beta_0 > 0$ (depending on α_0, δ_0 and the genus of S) such that:*

$$l_g(\{x \in c : \inf(i(g_x^l(\alpha_0), \lambda), i(g_x^r(\alpha_0), \lambda)) \leq \beta_0 l_g(\lambda)\}) \leq \delta_0 l_g(\lambda) + l_0.$$

Recall that $l_g(\lambda) = w l_g(c)$ if λ has weight w and support the simple closed curve c (see [9]).

The next proposition says that for points of c (on $\partial_+ C$) satisfying the reverse inequality (i.e. $\inf(n_l(x), n_r(x)) > \beta_0 l_g(c)\}$), the distance to the opposite boundary component of the convex core is near $\pi/2$. Cutting c by an orthogonal (totally geodesic) plane at such points, the assertion is about 2-dimensional AdS geometry.

Let C be a convex subset of AdS_2, with spacelike boundary, whose frontier in $AdS_2 \cup \partial_\infty AdS_2$ consists of 2 points, one on each component of $\partial_\infty AdS_2$. Let $\alpha_0 > 0$, x be a point of the upper boundary component $\partial_+ C$ of C, Π_x a support line of C at x, $P_{l,x}$ (resp. $P_{r,x}$) a support line of C at the point of $\partial_+ C$ at distance α_0 from the right (resp. the left) of x. We denote by $\phi_{r,x}$ (resp. $\phi_{l,x}$) the angle between Π_x and $P_{r,x}$ (resp. $P_{l,x}$). Let τ be a time-like geodesic orthogonal at x to Π_x.

Proposition 4. *For all $\epsilon > 0$, there exist $A > 0$, $\eta_0 > 0$ such that if*

$$\inf(\phi_{r,x}, \phi_{l,x}) \geq A$$

and

$$\alpha_0 \leq \eta_0$$

then

$$l(\tau \cap C) \geq \pi/2 - \epsilon.$$

Note that it is proved in [5] that we always have $l(\tau \cap C) < \pi/2$.

The next and last proposition applied in our situation enables us to compare the length of λ_+ (in case its support is a simple closed curve c) for m_+ and m_-, considering a totally geodesic time-like annulus containing c which is orthogonal, along c, to a support plane at some point of c.

Let M, m_+,m_-, λ_+,λ_- be as in Theorem 1.

Suppose that the support of λ_+ is a closed curve c. Let c_- be the curve obtained by intersecting the lower boundary component of the convex core of M with the totally geodesic timelike annulus orthogonal to a support plane at some point of c. Let K be a compact subset of $\mathcal{T}(S)$.

Proposition 5. *For all $\epsilon > 0$, there exists some $A > 0$ such that if $m_+ \in K$ and $l_{m_+}(\lambda_+) \geq A$ then $l_{m_-}(\lambda_+) \leq \epsilon l_{m_+}(\lambda_+)$.*

This last proposition extends to arbitrary λ_+ by density of weighted multicurves is $\mathcal{ML}(S)$, by continuity of length functions and continuity of m_- with respect to m_+ and λ_+.
Let us explain why the 3 propositions imply the theorem. In fact, the compactness theorem is a consequence of Proposition 5, which is a result of both Proposition 2 and 4.

Suppose there exist a sequence m_+^n in $\mathcal{T}(S)$ converging to $m_+^\infty \in \mathcal{T}(S)$ and a sequence ρ_l^n leaving any compact set in $\mathcal{T}(S)$. Then by Mess' diagram, the corresponding sequence of upper boundary's pleating lamination λ_+^n leaves any compact set in $\mathcal{ML}(S)$. In particular, its length, measured with respect to m_+^n, tends to infinity.

Let $\epsilon > 0$ and A be as in Proposition 5. There exists n_0 (depending on ϵ and A) such that for $n \geq n_0$, one has $l_{m_+^n}(\lambda_+^n) \geq A$, hence $l_{m_-^n}(\lambda_+^n) \leq \epsilon l_{m_+^n}(\lambda_+^n)$. Since m_+^n converges in $\mathcal{T}(S)$, this implies that m_-^n leaves every compact subset of $\mathcal{T}(S)$ (i.e tends to infinity), proving the properness of Φ.

4. Proof of the main propositions

We begin with the proof of Proposition 2, we need the following lemma.

Let S be a closed surface of negative Euler characteristic, $g \in \mathcal{T}(S)$, $\alpha_0 > 0$. Let c a closed geodesic for g. For any point $x \in c$, let $g_x^r(\alpha_0)$ (resp. $g_x^l(\alpha_0)$) be the geodesic segment of length α_0 (for the metric g), orthogonal at x to the right (resp. the left) of c. Let $n_l(x)$ (resp. $n_r(x)$) be the intersection number of $g_x^r(\alpha_0)$ (resp. $g_x^l(\alpha_0)$) with c, including x.

Lemma 6. *There exists a constant l_0 (depending only on α_0 and the genus of S) such that for all $\delta_0 > 0$, there exists some $\beta_0 > 0$ (depending on α_0, δ_0 and the genus of S) such that:*

$$l_g(\{x \in c : n_l(x) \leq \beta_0 l_g(c)\}) \leq \delta_0 l_g(c) + l_0.$$

The similar statement is true for $n_r(x)$ by the same argument. The proposition then follows by combining both statements.

Sublemma 7. *There exists γ_0 (depending on α_0) as follows. Let D_1, D_2 be two disjoint lines in the hyperbolic plane and let x be a point in the connected component of the complement in H^2 of $D_0 \cup D_1$ whose boundary contains those two lines. Suppose that $d(x, D_0) \leq \gamma_0$ and $d(x, D_1) \leq \gamma_0$. Then the geodesic segment of length α_0 starting orthogonally from D_0 and containing x intersects D_1.*



FIGURE 2. intersections de géodésiques hyperboliques planes

Proof of the sublemma. Let c_0 and c_1 be the geodesic segments joining x to its orthogonal projections on D_0 and D_1, respectively. Then $d(x, D_0) = l(c_0)$ and $d(x, D_1) = l(c_1)$. Let D be a hyperbolic line containing x and disjoint from D_0 and D_1, θ_0 and θ_1 be the respective angles between D and c_0, D and c_1. Then

$$\cosh(l(c_0)) \sin(\theta_0) = \cosh(d(D, D_0))$$

and

$$\cosh(l(c_1))\sin(\theta_1) = \cosh(d(D, D_1))$$

(See [13, page 88]). Since $d(D, D_0) \leq l(c_0) \leq \gamma_0$ and $d(D, D_1) \leq l(c_1) \leq \gamma_0$,

$$|\pi/2 - \theta_0| \leq \delta_0, |\pi/2 - \theta_1| \leq \delta_0$$

for sufficiently small γ_0 (depending on δ_0).

Let now c'_1 be the half-line extending c_1 on the other side of x. The angle between c_0 and c'_1 is less than $2\delta_0$. If γ_0 is small enough, then δ_0 is small and c'_1 intersects D_0 (which is orthogonal to c_0) at distance less than α_0. This is the required statement. □

Proof of the lemma. Let $\delta_0 > 0$, γ_0 as in the sublemma, and $0 < \beta_0 < \delta_0\gamma_0/(2\pi|\chi(S)|)$. Let $\overline{c} = \{x \in c : n_l(x) \leq \beta_0 l_g(c)\}$. Fix γ_0 as in the sublemma and consider the normal exponential map:

$$exp : \begin{array}{l} \overline{c} \times [0, \gamma_0] \to S \\ (s, r) \mapsto g^l_s(r) \end{array}$$

It is a distance increasing map so it increases areas. Moreover, the sublemma shows that each $x \in S$ has at most $n_0 + 1$ preimages in $\overline{c} \times [0, \gamma_0]$, where n_0 is the integer part of $\beta_0 l_g(c)$. Indeed, suppose that x is the image of both (y, r) and (y', r') and let $\overline{x}$,$\overline{y}$ and $\overline{y}'$ be lifts of x,y and y' to the universal cover H^2 of (S, g) chosen so that the corresponding lift of $\exp(\{y\} \times [0, r])$ (resp. $\exp(\{y'\} \times [0, r']))$ contains both $\overline{x}$ and $\overline{y}$ (resp. $\overline{x}$ and $\overline{y}$’). Then one of the following holds:

$exp(\{y\} \times [0, r])$ intersects the lift of c containing y'
$exp(\{y'\} \times [0, r'])$ intersects the lift of c containing y
$exp(\{y\} \times [0, \alpha_0])$ intersects the lift of c containing y' and conversely.

(The $3rd$ possibility is obtained by applying the SubLemma) This argument shows that there are at most n_0+1 preimages of x in $\overline{c}\times[0, \alpha_0]$.

Since the area of (S, g) is $2\pi|\chi(S)|$, it follows that

$$\gamma_0 l_g(\overline{c}) \leq 2\pi(n_0 + 1)|\chi(S)| \leq 2\pi(\beta_0 l_g(c) + 1)|\chi(S)| \leq \gamma_0(\delta_0 l_g(c) + l_0)$$

if one puts $l_0 = 2\pi|\chi(S)|/\gamma_0$. □

We now prove Proposition 4, we first need a lemma comparing the lengths of two past (resp. future) convex spacelike arcs, with the same endpoints, one being in the past (resp. future) of the other.

Lemma 8 (lengths of convex arcs with fixed endpoints)**.** *Let σ_0 and σ_1 be two future (resp. past) convex spacelike curves with the same* *endpoints in* AdS_2. *Suppose that* σ_0 *lies in the future (resp. past) of* σ_1. *Then*

$$l(\sigma_0) \geq l(\sigma_1).$$

Proof of the lemma. We first examine the case where σ_0 and σ_1 are piecewise geodesic arcs.

First, suppose σ_0 is a geodesic. Then by induction on the number of geodesic arcs in σ_1, this case is a consequence of Sublemma 4.6 of [5] (the "reverse triangle inequality").

Now suppose σ_0 is a piecewise geodesic arc. Consider the geodesic arc σ_0' joining the endpoints of σ_0. By orthogonal projection of each vertex of σ_1 on σ_0', we get timelike arcs joining those vertices to points of σ_0' and orthogonal to σ_0'. Now we can pull back σ_1 along those arcs (with the normal exponential map) until it reaches (at least) one point of σ_0. We get a piecewise geodesic arc σ_1', which is longer than σ_1, spacelike and future convex but this time σ_1' and σ_0 have a common point. We can then apply the induction argument on both sides of this common point on σ_0 and σ_1'.

Then the general case follows by approximation. □

We keep the same notations as in the statement of Proposition 4 .

Proof of Proposition 4. Letting x_l (resp. x_r) be the intersection point of Π_x and $P_{l,x}$ (resp. $P_{r,x}$), we note that the (spacelike) distance between x and x_l (resp. x and x_r) is less than α_0, by Lemma 1.

The configuration made by the three spacelike lines and the spacelike line Π' joining the two points at infinity of $P_{l,x}$ and $P_{r,x}$ in the past of Π_x admits a well-known limit situation: when α_0 goes to zero, x_l and x_r tend to x, moreover when $\phi_{l,x}$ and $\phi_{r,x}$ tend to infinity, $P_{l,x}$ and $P_{r,x}$ tend to lightlike lines so as to let Π' tend to the dual line Π^* to x, at (timelike) distance $\pi/2$ from x.

The result follows since by convexity,

$$l(\tau \cap C) \geq l(I^+(\Pi') \cap \tau)$$

□

Note that in our case, since α_0 tends to 0, having our intersection numbers $n_l(x)$ and $n_r(x)$ tend to ∞ is equivalent to having the angles $\phi_{l,x}$ and $\phi_{r,x}$ tend to ∞ (since the left and right geodesic segments of length α_0 at x meet c with an angle near from $\pi/2$)

It remains to prove Proposition 5 . We keep the same notations as in the statement of this proposition.



FIGURE 3. cutting the convex core by a plane orthogonal to a support line

Proof of Proposition 5. Let $\epsilon > 0$, let K be a compact subset of $\mathcal{T}(S)$. We choose $\epsilon' > 0$ (say $\epsilon' < \pi/2$), such that

$$\cos(\pi/2 - \epsilon') \leq \epsilon/2.$$

Then there are constants A' and η_0 such that if

$$\inf(\phi_{r,x}, \phi_{l,x}) \geq A'$$

and

$$\alpha_0 \leq \eta_0$$

then

$$l(\tau \cap C) \geq \pi/2 - \epsilon'$$

with the notations of Proposition 4 .

Let $A(c)$ be a totally geodesic timelike annulus orthogonal to a support plane of C along c. Let $c_- = A(c) \cap \partial_- C$ and let c'_- be the curve



FIGURE 4. dual lines

at distance $\pi/2-\epsilon'$ in the past of c (lying in $A(c)$). Then c'_- is a future convex curve and

$$l(c'_-) = \cos(\pi/2-\epsilon')l_{m_+}(c),$$

hence

$$l(c'_-) \leq (\epsilon/2)l_{m_+}(c).$$

Consider now the orthogonal projection Pr_- from c_- to c'_-. There are two types of points on c_-, those at distance greater than or equal to $\pi/2-\epsilon'$ from c and the others.

Let β_0 as in Proposition 2 and $\delta_0 = \epsilon/4$. Proposition 4 gives the existence of constants A' and η_0 (where $\alpha_0 \leq \inf(\eta_0, \epsilon/2)$), we choose c such that $l_{m_+}(c) \geq A'/\beta_0$. Let

$$B_- = \{\tau_x \cap \partial_- C, x \in c, \inf(n_l(x), n_r(x)) \geq \beta_0 l(c)\},$$

then points of B_- belongs to

$$\{y \in c_-, d(y,c) \geq \pi/2-\epsilon'\}.$$

The points of the first type are in the past of c'_-, so by Lemma 1,

$$l(B_-) \leq l(Pr_-(B_-)),$$

hence by composition of projections,

$$l(B_-) \leq (\epsilon/2)l(Pr(B_-)),$$

where Pr is the projection from c_- to c.
Let

$$B = \{\tau_x \cap \partial_- C, x \in c, \inf(n_l(x), n_r(x)) \leq \beta_0 l(c)\}.$$



FIGURE 5. comparing length on top and bottom of the convex core

Lemma 1 gives

$$l(\{x \in c, \inf(n_l(x), n_r(x)) \leq \beta_0 l(c)\}) \geq l(B)$$

Then, by choosing $A \geq \sup(A'/\beta_0, 4l_0/\epsilon)$ (where l_0 comes from Proposition 4),

$$l(\{x \in c, \inf(n_l(x), n_r(x)) \leq \beta_0 l(c)\}) \leq \delta_0 l(c) + l_0 \leq \epsilon l(c)/4 + \epsilon l(c)/4$$

hence

$$\epsilon l(c)/2 \geq l(B) \quad .$$

We get the desired result by addition, then by passing to the limit for a general lamination. □

5. COMPUTING DEGREES

In this section, we investigate the degree of the proper map Φ. Our main aim is the

Proposition 9 (degree one theorem)**.** *Let Φ be the map defined in section 2. Then Φ has degree one.*

Recall that the degree of proper maps between manifolds is well defined and that a map whose degree is nonzero is surjective. Thus the last proposition ends the proof of the main theorem of the article.

In [2], they prove the existence of a map Φ_{k_+,k_-} from $\mathcal{G}H(S)$ (GHMC AdS_3 structures with Cauchy surface homeomorphic to S, identified with $\mathcal{T}(S)\times\mathcal{T}(S)$) to $\mathcal{T}(S)\times\mathcal{T}(S)$ which associates to a GHMC AdS3 manifold the pair of conformal structures of the unique past convex (resp. future convex) surfaces with constant sectional curvatures equal to k_+ and k_- respectively (see also [4] where this map is considered).

They also prove that the map φ

$$(k^+,k^-,\rho_l,\rho_r)\mapsto\Phi_{k_+,k_-}(\rho_l,\rho_r)=(\Phi^+_{k_+,k_-}(\rho_l,\rho_r),\Phi^-_{k_+,k_-}(\rho_l,\rho_r))$$

is continuous (lemma 12.4) on $(-\infty,-1)^2\times\mathcal{T}(S)\times\mathcal{T}(S)$.

The continuity on $(-\infty,-1]^2\times\mathcal{T}(S)\times\mathcal{T}(S)$ follows from the following lemma:

Lemma 10 (convergence of a sequence of hyperbolic metrics)**.** *Let m be a point of $\mathcal{T}(S)$ and C_n be a sequence of real numbers greater than one which converge to* 1*.*

Then any sequence $m_n\in\mathcal{T}(S)$ satisfying

$$m_n\leq C_n m$$

converge to m.

This is a consequence of a classical fact proved by Thurston [14]. Indeed, by the following lemma, such a sequence admits a convergent subsequence, whose limit m_∞ satisfies $m_\infty\leq m$. Then Thurston proved that this gives $m_\infty=m$.

The map φ, when restricted to any set of the form $[-C,-1]^2\times\mathcal{T}(S)\times\mathcal{T}(S)$ (for any $C>1$), is proper thanks to the following lemmas (which are corollaries of results from [14]):

Lemma 11 (compactness of sets of metrics first version)**.** *Let m be a point of $\mathcal{T}(S)$ and let $C>1$.*

Then the set of metrics m' in $\mathcal{T}(S)$ such that: $Cm'\geq m$ is compact in $\mathcal{T}(S)$.

The same lemma is true when considering metrics m' such that $m'\leq Cm$.

By a slight extention we get the following lemma:

Lemma 12 (compactness of sets of metrics revisited)**.** *Let K be a compact of $\mathcal{T}(S)$, let $C>1$. Then the set of metrics m' such that $Cm'>m$ for some $m\in K$ is compact.*

Consider the slicing of an AdS GHMC manifold given by the map φ . For any $k < -1$, let k^* be the curvature of the dual to the surface of curvature k. Then we have:(formula for $k*$)

Let us then prove the properness of the map φ. Let (m_n^+) and (m_n^-) be convergent sequences in $\mathcal{T}(S)$ and ρ_n^l, ρ_n^r, k_n^+ and k_n^- be sequences such that the image of

$$(k_n^+, k_n^-, \rho_n^l, \rho_n^r)$$

by φ is $((m_n^+), (m_n^-))$.

By properness of $\Phi_{-1,-1} = \Phi$ (proved in the previous section) and the fact that

$$C\Phi^+_{k_+,k_-} < \Phi^+_{-1,-1}$$

and

$$C\Phi^-_{k_+,k_-} < \Phi^-_{-1,-1},$$

the sequences k_n^+,k_n^-,ρ_n^l,ρ_n^r stay in a compact set, hence they admit convergent subsequences.

By invariance of the degree of a map under a proper homotopy, all the maps Φ_{k_+,k_-} have the same degree, which is given by this last lemma:

Lemma 13 (degree of Φ_{k,k^*})**.** *The map* Φ_{k,k^*} *has degree one*

In fact it is a homeomorphism.

References

[1] Lars Andersson, Thierry Barbot, Riccardo Benedetti, Francesco Bonsante, William M. Goldman, François Labourie, Kevin P. Scannell, and Jean-Marc Schlenker. Notes on: "Lorentz spacetimes of constant curvature" [Geom. Dedicata **126** (2007), 3–45; mr2328921] by G. Mess. *Geom. Dedicata*, 126:47–70, 2007.

[2] Thierry Barbot, François Béguin, and Abdelghani Zeghib. Prescribing Gauss curvature of surfaces in 3-dimensional spacetimes: application to the Minkowski problem in the Minkowski space. *Ann. Inst. Fourier (Grenoble)*, 61(2):511–591, 2011.

[3] Lipman Bers. Simultaneous uniformization. *Bull. Amer. Math. Soc.*, 66:94–97, 1960.

[4] F. Bonsante, G. Mondello, and J.-M. Schlenker. A cyclic extension of the earthquake flow. *ArXiv e-prints.* To appear*, Geometry and Topology*, June 2011.

[5] Francesco Bonsante and Jean-Marc Schlenker. Fixed points of compositions of earthquakes. *Duke Math. J.*, 161(6):1011–1054, 2012.

[6] R. D. Canary, D. B. A. Epstein, and P. L. Green. Notes on notes of Thurston [mr0903850]. In *Fundamentals of hyperbolic geometry: selected expositions*, volume 328 of *London Math. Soc. Lecture Note Ser.*, pages 1–115. Cambridge Univ. Press, Cambridge, 2006. With a new foreword by Canary.

[7] D. B. A. Epstein and A. Marden. Convex hulls in hyperbolic space, a theorem of Sullivan, and measured pleated surfaces. In *Analytical and geometric aspects of hyperbolic space (Coventry/Durham, 1984)*, volume 111 of *London Math. Soc. Lecture Note Ser.*, pages 113–253. Cambridge Univ. Press, Cambridge, 1987.

[8] D. B. A. Epstein and V. Markovic. The logarithmic spiral: a counterexample to the $K = 2$ conjecture. *Ann. of Math. (2)*, 161(2):925–957, 2005.

[9] Steven P. Kerckhoff. Earthquakes are analytic. *Comment. Math. Helv.*, 60(1):17–30, 1985.

[10] François Labourie. Métriques prescrites sur le bord des variétés hyperboliques de dimension 3. *J. Differential Geom.*, 35(3):609–626, 1992.

[11] François Labourie and Jean-Marc Schlenker. Surfaces convexes fuchsiennes dans les espaces lorentziens à courbure constante. *Math. Ann.*, 316(3):465–483, 2000.

[12] Geoffrey Mess. Lorentz spacetimes of constant curvature. *Geom. Dedicata*, 126:3–45, 2007.

[13] John G. Ratcliffe. *Foundations of hyperbolic manifolds*, volume 149 of *Graduate Texts in Mathematics*. Springer, New York, second edition, 2006.

[14] W. P. Thurston. Minimal stretch maps between hyperbolic surfaces. *ArXiv Mathematics e-prints*, January 1998.

[15] William P. Thurston. Three-dimensional geometry and topology. Originally notes of lectures at Princeton University, 1979. Recent version available on http://www.msri.org/publications/books/gt3m/, 1980.

[16] William P. Thurston. Earthquakes in two-dimensional hyperbolic geometry. In *Low-dimensional topology and Kleinian groups (Coventry/Durham, 1984)*, volume 112 of *London Math. Soc. Lecture Note Ser.*, pages 91–112. Cambridge Univ. Press, Cambridge, 1986.

Institut de Mathématiques de Toulouse, UMR CNRS 5219, Université Toulouse 3, 118 route de Narbonne, 31062 Toulouse cedex 9, France

E-mail address: `boubacar.diallo.math@gmail.com`